

hydrogens in the alkylene chain may optionally be substituted by any one of C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; or
one or more of the methylene groups may optionally be substituted by a heteroatom selected from O, N or S;

R¹ is selected from hydrogen, a branched or straight C₁-C₆ alkyl, C₂-C₆ alkenyl, C₃-C₈ cycloalkyl, C₄-C₈ (alkyl-cycloalkyl) wherein the alkyl is a C₁-C₂ alkyl and the cycloalkyl is a C₃-C₆ cycloalkyl;

R² is selected from any of:

- P₁
- (i) hydrogen;
 - (ii) a straight or branched C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
 - (iii) -[(CH₂)_q-aryl], wherein the aryl may optionally be substituted by 1 or 2 substituents Y, wherein each Y is as defined below; and wherein q is an integer from 0 to 3;
 - (iv) -[(CH₂)_r-heteroaryl] wherein the heteroaryl has from 5 to 10 atoms, each heteroatom being selected from any of S, N and O and wherein the heteroaryl may be substituted by 1 or 2 substituents Y, wherein each Y is as defined below; and wherein r is an integer from 0 to 3;
 - (v) C₃-C₁₀ cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryls, where each heteroaryl has from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (vi) C₆-C₁₀ aryl, optionally and independently substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (vii) a heteroaryl having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; wherein the aryl and heteroaryl may optionally and

independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;

or R¹ and R² may optionally form a heterocyclic ring;

R³ is selected from any one of:

- D₁*
- (i) hydrogen;
 - (ii) a straight or branched C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
 - (iii) -[(CH₂)_q-aryl] wherein q is an integer from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (iv) a heteroaryl-(C₅-C₁₀alkyl), wherein the heteroaryl has from 5 to 10 atoms, each heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (v) a C₃-C₁₀ cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (vi) -[(C₃-C₆ cycloalkyl)-(CH₂)_q] wherein q is an integer from 1 to 3;

R⁴ is selected from:


- (i) hydrogen;
- (ii) a straight or branched C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
- (iii) -[(CH₂)_q-aryl] wherein q is an integer from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;

- 21
- (iv) heteroaryl-(C₅-C₁₀ alkyl), wherein the heteroaryl has from 5 to 10 atoms, each heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (v) a C₃-C₁₀ cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O;
and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (vi) a C₆-C₁₀ aryl, optionally and independently substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
 - (vii) a heteroaryl having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein Y is as defined below;

R⁵ is selected from:

- (i) hydrogen;
- (ii) a straight or branched C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
- (iii) -(CH₂)_q-aryl] wherein q is an integer from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
- (iv) a heteroaryl-(C₅-C₁₀ alkyl), wherein the heteroaryl has from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted 1 or 2 substituents Y, wherein each Y is as defined below;

- (v) a C₃-C₁₀ cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;



wherein R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each and independently selected from:

- (a) hydrogen;
- (b) a straight or branched C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
- (c) -[(CH₂)_q-aryl] wherein q is an integer from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of the S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;
- (d) a heteroaryl-(C₅-C₁₀ alkyl), wherein the heteroaryl has from 5 to 10 atoms, each heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted 1 or 2 substituents Y, wherein each Y is as defined below;
- (e) a C₃-C₁₀ cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted 1 or 2 substituents Y, wherein each Y is as defined below;
- (f) a C₆-C₁₀ aryl, optionally and independently substituted by one or more heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O, and wherein the heteroaryl may optionally and

independently be substituted by 1 or 2 substituents Y, wherein each Y is as defined below;

or R⁴ and R⁵ may optionally form a heterocyclic ring;

Y is each and independently selected from any of: hydrogen, CH₃; -(CH₂)_{p1}CF₃; halogen; C₁-C₃ alkoxy; hydroxy; -NO₂; -OCF₃; -CONR^aR^b; -COOR^a; -COR^a; -(CH₂)_{p2}NR^aR^b; -(CH₂)_{p3}CH₃; (CH₂)_{p4}SOR^aR^b; -(CH₂)_{p5}SO₂R^a; -(CH₂)_{p6}SO₂NR^a; C₄-C₈(alkyl-cycloalkyl) wherein the alkyl is a C₁-C₂ alkyl, and the cycloalkyl is a C₃-C₆ cycloalkyl; 1 or 2 heteroaryls having from 5 to 10 atoms, each heteroatom being selected from any of S, N and O; and oxides selected from N-oxides or sulfoxides; and wherein:

R^a and R^b are each and independently selected from hydrogen, a branched or straight C₁-C₆ alkyl, a C₁-C₆ alkenyl, a C₃-C₈ cycloalkyl; and wherein:

p¹, p², p³, p⁴, p⁵ and p⁶ are each and independently 0, 1 or 2;

as well as pharmaceutically acceptable salts, isomers, hydrates, and isoforms thereof,

with the proviso that when R¹=R³=R⁴=R⁵=H, then R² is not hydrogen or a straight or branched C₁-C₆ alkyl and when R²=R³=R⁴=R⁵=H then R¹ is not hydrogen or a straight or branched C₁-C₆ alkyl.

40. (Once amended) A compound wherein said compound is:

